

**On the Kraichnan Model of Passive Scalar Advection
Near the Batchelor limit**

Alain Pumir
Institut Non Linéaire de Nice, 1361 route des Lucioles
F-06560, Valbonne, France

Boris I. Shraiman
Bell Laboratories, Lucent Technologies
700 Mountain Avenue, Murray Hill, NJ 07974

&

Eric D. Siggia
Cornell University, Laboratory of Atomic & Solid State Physics
Ithaca, NY 14853-2501

Abstract

The third order correlation function of the scalar field advected by a Gaussian random velocity, with a spatial scaling exponent $2 - \epsilon$, and in the presence of a mean gradient, is calculated perturbatively in $\epsilon \ll 1$. This expansion corresponds to the regime close to Batchelor's advection by linear diffeomorphisms. The scaling exponent is found to be equal to 1 in dimensions 2 and 3, up to corrections smaller than $\mathcal{O}(\epsilon)$, implying an anomalous scaling of the third order correlation function and the persistence of small scale anisotropy.

PACS numbers :

47.27Gs. - Isotropic turbulence; homogeneous turbulence.

47.27Te. - Convection and heat transfer.

The investigation of the statistics of the passive scalar field advected by random flow is interesting for the insight it offers into the origin of intermittency and anomalous scaling of turbulent fluctuations. The problem studied in this paper is stated simply by:

$$\partial_t \Theta + (\vec{u} \cdot \nabla) \Theta = \kappa \nabla^2 \Theta \quad (1)$$

with the scalar field Θ forced by the externally imposed gradient, g . It is convenient to subtract out the gradient and study the fluctuating field, $\theta(r) = \Theta(r) - gr$. It turns out that even a Gaussian random, but scale invariant, velocity field results in non-trivial anomalous scaling of the passive scalar structure function, $\langle (\Theta(r) - \Theta(0))^n \rangle$ for $n > 2$. This has been argued by Kraichnan¹, via a plausible closure scheme, for a "white noise" model where :

$$\langle v_a(r, t) v_b(r', t') \rangle = \delta(t - t') C_{ab}(r - r') \quad (2a)$$

with :

$$D_{ab}(r) = C_{ab}(0) - C_{ab}(r) = D_0 \left((d - 1 + \zeta_v) \delta_{ab} - \zeta_v \frac{r^a r^b}{|r|^2} \right) |r|^{\zeta_v} \quad (2b)$$

(where ζ_v is the scaling exponent and d the space dimension) which he has introduced some 30 years ago.² The existence of an anomalous scaling has been demonstrated explicitly by Gawedski and Kupiainen³, and Chertkov et al.⁴ for certain limits of this Kraichnan model and by Shraiman and Siggia⁵ for a generalized phenomenological model where temporal correlation of the advecting field is set by eddy turnover. These calculations are based on the so called Hopf equations - the stationarity conditions of the equal-time multipoint correlators. For the white velocity case these can be derived exactly^{6,7} extending the original analysis of the 2-point function by Kraichnan⁸. They have the form:

$$\begin{aligned}
& \sum_{i \neq j}^N \left(D_{ab}(r_i - r_j) + \kappa \delta_{ab} \right) \partial_{r_i}^a \partial_{r_j}^b < \theta(r_1) \dots \theta(r_N) > \\
& = \sum_{i \neq j}^N g_a g_b C_{ab}(r_i - r_j) < \theta \dots >_{ij}^{N-2} \\
& \quad - 2 \sum_{i \neq j}^N g_a D_{ab}(r_i - r_j) \partial_j^b < \theta \dots >_i^{N-1}
\end{aligned} \tag{2c}$$

(with implicit summation over repeating indices). We restrict ourselves to the inertial range of scales, where r is large enough so that the molecular diffusivity can be neglected: $r \gg \eta \equiv (\kappa/D_0)^{1/\zeta_v}$.

The analysis of Ref.3 is based on the expansion of Eq.(2) in $\zeta_v \ll 1$ about the diffusion limit $\zeta_v = 0$, while we consider the complementary limit of $\zeta_v = 2 - \epsilon$, $\epsilon \ll 1$. Reality for the white velocity model, $\zeta_v = 4/3$, lies inbetween. The expansion in small ϵ is more involved than what was required in Refs 3,4 for two reasons. There are an infinite number of degenerate modes for $\epsilon = 0$ which are all mixed by the perturbation, which itself is singular^{5,9}. That is the perturbation is formally small because of ϵ , but in certain restricted regions of configuration space it is the biggest term in the equation. It must be treated by the method of matched asymptotic expansions. The exponent we find for the third order correlator^{10,11} $\lambda_3 \approx 1$ implies that the anisotropy introduced by the mean gradient, g on the large scales, decays more slowly as one descends in scale than that predicted by K41 theory^{12,13} (which for $\zeta_v = 2 - \epsilon$ predicts an exponent $1 + \epsilon$). Since the experimental exponent is also approximately one¹⁴, it will be of interest to compare also the full coordinate dependence of the three point correlation function when the latter becomes available from experiment or simulations. One way of expressing our result for this correlation function, is as an expansion in the degenerate modes of the $\epsilon = 0$ problem. Our matching determines all the coefficients explicitly.

Determination of the anomalous exponents reduces to finding the zero modes of the linear operator entering the Hopf equation,^{3,4,5} i.e. the left hand side of Eq.(2) which

in the present model (and $\kappa \rightarrow 0$ limit) is the generalized Richardson diffusion operator: $L(d, \zeta_v) \equiv \sum_{i \neq j}^3 D_{ab}(r_i - r_j) \partial_{r_i}^a \partial_{r_j}^b$. The $\zeta_v = 2$ case is the Batchelor limit¹⁵ which is constrained by an overall $SL(2) \times SO(d)$ symmetry so that the spectrum of $L_0 \equiv L(d, 2)$, also referred to here as the Batchelor-Kraichnan operator, can be completely constructed with the help of Lie algebraic methods^{5,16}. Here it will serve as a starting point for the perturbation theory for $\zeta_v = 2 - \epsilon$:

$$L(d, 2 - \epsilon) = L_0(d) - \epsilon L_1(d) \quad (3)$$

to the leading order in $\epsilon \ll 1$. (Note, that we are ultimately interested in the physical case of $\zeta_v = 4/3$.) The perturbation expansion around $\zeta_v = 2$ is a singular problem which however can be addressed by the method introduced in Ref.9, as we explain now.

Let us start with L_0 . It is convenient to introduce the variables $\vec{\rho}_1 = (\vec{r}_1 - \vec{r}_2)/\sqrt{2}$ and $\vec{\rho}_2 = (\vec{r}_1 + \vec{r}_2 - 2\vec{r}_3)/\sqrt{6}$ (On occasion we shall refer to $i = 1, 2$ -index labelling the ρ vectors as the "pseudo-space" index to distinguish it from the d -dimensional real space.) Next we "factorize": $\rho_i^a = \sum_{i'} R_{ii'}(\chi) \xi_{i'} \eta_{i'}^a$, where R represents pseudo-space rotations by χ , and $\hat{\eta}_{1,2}$ are two orthogonal unit vectors. In $d = 3$, we also define $\hat{\eta}_3 \equiv \hat{\eta}_1 \wedge \hat{\eta}_2 = \vec{\rho}_1 \wedge \vec{\rho}_2 / |\vec{\rho}_1 \wedge \vec{\rho}_2|$ each component of which is invariant under the action of $SL(2)$ ⁹. Another important invariant is the area of the $\vec{r}_1, \vec{r}_2, \vec{r}_3$ triangle: $\zeta \equiv |\vec{\rho}_1 \wedge \vec{\rho}_2| = \xi_1 \xi_2$.

The zero modes of $L_0(d)$ for $d = 2, 3$ have been constructed in Ref. 5,9; e.g. in $d = 3$, the complete set of eigenfunctions has the form:

$$\psi_{\nu, q, l, m, m'}^\lambda = e^{iq\chi} \zeta^{\frac{\lambda}{2}} P_\nu^{q, m'}(\xi) D_{m, m'}^l(\hat{\eta}) \quad (4),$$

where $\xi \equiv (\xi_1^2 + \xi_2^2)/2\xi_1\xi_2$, $D_{m, m'}^l(\hat{\eta})$ is the matrix element of the representation of the $SO(3)$ group¹⁷ of order l and $P_\nu^{q, m'}$ is the Jacobi function.¹⁸ (Note that quantum number m' corresponds to rotations of the $\hat{\eta}$ triad about $\hat{\eta}_3$ in pseudo-space.) To ensure analyticity in the $\zeta = |\rho_1 \wedge \rho_2| \rightarrow 0$ limit (which corresponds to all 3 points of the correlator being on one line), $\lambda/2 - \max(\nu, -\nu - 1)$ must be a positive integer. This is because as $\zeta \rightarrow 0$, $\xi \sim \zeta^{-1} \rightarrow \infty$ and $P_\nu^{q, m}(\xi) \sim \xi^{\max(\nu, -\nu - 1)}$.

The 3-order structure function, or the skewness, which is the physical object of interest has odd spatial parity and hence is only non-zero in as much as the mean scalar gradient, g , introduces a particular direction. Hence the relevant eigenfunctions are the p-waves, $l = 1$. The zero mode of $L_0(d)$ corresponding to the smallest exponent λ is obtained for $\frac{\lambda}{2} = \nu$ yielding, in the $l = 1$ sector, $\lambda = 1$ independent of d .

We shall need the explicit form of the L_0 operator:

$$\frac{1}{2d} L_0 \psi_{l=1}^\lambda(w, \chi, \hat{\eta}) = \partial_\xi((\xi^2 - 1)\partial_\xi \psi) + \frac{(\partial_\chi^2 - I_3^2) - 2i\xi I_3 \partial_\chi}{4(\xi^2 - 1)} \psi - \nu(\nu + 1)\psi \quad (5),$$

where $\nu(\nu + 1) \equiv \frac{d-2}{2d}(\frac{\lambda^2}{2d} + \lambda) - \frac{d+1}{2d}l(l+d-2)$ and $I_3 \equiv \frac{1}{i}(\eta_1 \partial_2 - \eta_2 \partial_1)$. In agreement with Eq.(5), the ∂_χ and I_3^2 are diagonalized by $\exp(iq\chi)$ and $\hat{\eta}_1 \pm i\hat{\eta}_2$ the latter corresponding to the $l = 1, m' = \pm 1$ sector. Requiring the left hand side of Eq.(5) to vanish would make it into a Legendre equation¹⁸ with ν and hence λ entering as an eigenvalue.

Next we define the perturbation operator in Eq.(3):

$$L_1(d) = \mathcal{L} + \frac{d-1}{2d} (l(l+d-2) + \lambda^2 - d\lambda - \frac{1}{d-1} L_0) \quad (6a),$$

with :

$$\mathcal{L} \equiv \sum_{\mathcal{S}_3} -\ln(|\rho_1|) \times \left((d+1)\rho_1^2(\partial_1^a \partial_1^a - \frac{1}{3}\partial_2^a \partial_2^a) - 2\rho_1^a \rho_1^b (\partial_1^a \partial_1^b - \frac{1}{3}\partial_2^a \partial_2^b) \right) \quad (6b).$$

In Eq.(6b), the summation extends over all the cyclic permutations of $(\vec{r}_1, \vec{r}_2, \vec{r}_3)$, resulting in the following symmetry for the ρ : $\rho_1 \rightarrow -\frac{\rho_1}{2} \pm \frac{\sqrt{3}}{2}\rho_2$ and $\rho_2 \rightarrow \mp \frac{\sqrt{3}}{2}\rho_1 - \frac{\rho_2}{2}$. The expansion breaks down for $-\ln|\rho_1| < \epsilon^{-1}$. When $\xi \rightarrow \infty$, *i.e.*, when the three points $\vec{r}_1$, $\vec{r}_2$ and $\vec{r}_3$ are almost aligned, the operator $\epsilon\mathcal{L}$ becomes much larger than the Batchelor-Kraichnan operator L_0 . This can be seen by expanding the full operator in the limit $\xi \rightarrow \infty$. Defining $\psi_{l=1}^\lambda \equiv \zeta^{\frac{\lambda}{2}} \varphi(\xi, \eta, \chi)$, one finds, for $d=3$ and the $l = 1$ sector:

$$\begin{aligned} \mathcal{L}\varphi = & -\ln[1 - (1 - \xi^{-2}) \cos(2\chi)] \times [\xi^2 \mathcal{L}_2 \varphi + \xi \mathcal{L}_1 \varphi + \mathcal{L}_0 \varphi + \mathcal{O}(1/\xi)] \\ & + (\chi \rightarrow \chi + \frac{2\pi}{3}) + (\chi \rightarrow \chi - \frac{2\pi}{3}) \end{aligned} \quad (7a).$$

The operators $\mathcal{L}_i$ are :

$$\mathcal{L}_2\varphi = \frac{4}{3}(\cos^2\chi - 1)(4\cos^2\chi - 1)\left(-(2\xi\partial_\xi - \lambda)^2 + 4\eta_1\partial_{\eta_1}\right)\varphi \quad (7b),$$

$$\mathcal{L}_1\varphi = \frac{32}{3}\cos\chi\sin\chi(\cos^2\chi - 1)\left((2\xi\partial_\xi - 1)(\eta_2\partial_{\eta_1} - \eta_1\partial_{\eta_2}) - 2\eta_2\partial_{\eta_1}\right)\varphi \quad (7c),$$

and :

$$\begin{aligned} \mathcal{L}_0\varphi = & -\frac{2}{3}(\cos^2\chi - 1)(4\cos^2\chi - 1)(\partial_{\chi^2} + 5)\varphi \\ & + \frac{64}{3}\sin^3\chi\cos\chi\partial_\chi\varphi + \frac{4}{3}(1 + 4\cos^2\chi - 8\cos^4\chi)\eta_1\partial_{\eta_1}\varphi \end{aligned} \quad (7d).$$

In Eq.(7d), λ has been set equal to 1 - its unperturbed value - since the corrections would be higher order in ϵ . The singular nature of the perturbation follows from the fact that the $\mathcal{L}_2$ term enters with prefactor ξ^2 so that when $\xi \gg (1/\epsilon)^{1/2}$, $\epsilon\mathcal{L} \gg L_0$. This situation calls for the "boundary layer" type matched asymptotic analysis which we outline below.

Let us assume $\lambda = 1 + \epsilon\delta$, define the rescaled 'inner' variable $z = \epsilon^{1/2}\xi$, and introduce the function $\varphi(z, \chi, \eta) = z^{\frac{\lambda}{2}}(\phi_1(z, \chi)\hat{\eta}_1 + i\phi_2(z, \chi)\hat{\eta}_2)$. The prefactor is chosen to offset the scaling factor $\zeta^{\lambda/2}$ (see Eq.(4)) which vanishes for collinear points. Physics requires that ϕ_i is bounded when $z \rightarrow \infty$. With this change of variable and functions, the problem can be written, provided $\chi \gtrsim \xi^{-1}$ as :

$$\left(\left((z^2\partial_z^2 + 3z\partial_z) + \frac{4}{9}z^2U(\chi)((z\partial_z)^2 - \eta_1\partial_{\eta_1})\right) + \epsilon^{\frac{1}{2}}\hat{L}_1 + \epsilon\hat{L}_2 + \dots\right)(\phi_1\eta_1 + i\phi_2\eta_2) = 0 \quad (8a),$$

with

$$\begin{aligned} U(\chi) = & \left((\cos^2\chi - 1)(4\cos^2\chi - 1)\ln(1 - \cos(2\chi))\right. \\ & \left.+ (\chi \rightarrow \chi + 2\pi/3) + (\chi \rightarrow \chi - 2\pi/3)\right) \end{aligned} \quad (8b).$$

The operators $\hat{L}_1$ and $\hat{L}_2$ can be deduced from a systematic expansion of the operator in powers of ϵ starting from Eq.(7a-d).

The boundary conditions at infinity imply that when $z \rightarrow \infty$, the solution is a function of χ only. By direct substitution, one finds that $\phi_1 = 0$ and $\phi_2 = a(\chi)$, where $a(\chi)$ is an unknown function, decomposed for convenience as a Fourier series in χ : $a(\chi) = \sum_q \hat{a}_q e^{iq\chi}$. When $z \rightarrow 0$, the problem reduces to the unperturbed Batchelor-Kraichnan operator, up

to small corrections. In the matching region, defined by $z \rightarrow 0$ but $\xi \rightarrow \infty$, or equivalently, $1 \ll \xi \ll \epsilon^{-1/2}$, the ξ dependence of each Fourier mode in χ , q , must match with the asymptotic behavior of the eigenmodes of L_0 (Eq.(4,5)) which is best found via their integral representation given in Ref.5,9,17. One finds that the functions $\phi_{i,q}$ must behave as: $\phi_{1,q} = \frac{\epsilon^{1/2}|q|}{2z} + \frac{\epsilon(1-q^2)}{4z^2} + \dots$ and $\phi_{2,q} = (1 - \epsilon \frac{q^2}{8z^2} + \dots) \text{sign}(q)$. The crossover equation, Eq.(8), can be solved analytically, to the leading order in ϵ , and the imposition of the matching conditions determines $a(\chi)$ via an eigenvalue equation for δ :

$$U(\chi)(\partial_\chi^2 + 1)a(\chi) + 6\delta a = 0 \quad (9).$$

The analysis in $d = 2$ can be carried out in a completely similar way. As in the 3-dimensional case, the behavior for $\xi \rightarrow \infty$ is of the form $\phi = a_{2d}(\chi)\xi^{\lambda/2} + \mathcal{O}(\xi^{\lambda/2-1})$, and the function a_{2d} is determined by a matching condition. Surprisingly, the equation determining $a_{2d}(\chi)$, and the correction to the scaling exponent, $\delta = (\lambda - 1)/\epsilon$ is identical to Eq.(9).

Before solving Eq.(9), one needs to determine the appropriate boundary conditions. Because the 3-point correlation function must be odd under $\vec{\rho}_i \rightarrow -\vec{\rho}_i$, implying that $a(\chi + \pi) = -a(\chi)$. This, together with the periodicity $a(\chi + 2\pi/3) = a(\chi)$, resulting from the invariance under cyclic permutation of $\vec{r}_1$, $\vec{r}_2$ and $\vec{r}_3$, implies that $a(\chi + \pi/3) = -a(\chi)$. The limit $\chi \rightarrow 0$ corresponds to the case where $\vec{r}_1$ and $\vec{r}_2$ come close together : $|\vec{r}_2 - \vec{r}_1| \ll |\vec{r}_3 - \vec{r}_1|, |\vec{r}_2 - \vec{r}_1|$. In this limit, the correlation function must be invariant when $\vec{r}_1$ and $\vec{r}_2$ are permuted, implying that $a(\chi)$ must be even. Since a is even near $\chi = 0$ and a is antiperiodic with period $\pi/3$, $a(\pi/6) = 0$.

At small, but finite ϵ , Eq.(9) reduces for $\chi \rightarrow 0$, to :

$$-\chi^2 \ln(\chi) a''(\chi) + \delta a(\chi) = 0 \quad (10a).$$

Introducing the change of variables : $y \equiv -\ln(\chi)$ and $f(y) \equiv a(\chi)$, Eq.(10a) reduces to the following (Kummer) equation :

$$y(f'' + f') + \delta f = 0 \quad (10b).$$

The behavior of the solution when $\chi \rightarrow 0$ is:

$$f(y) \sim y^{-\delta} \quad \text{when} \quad y \rightarrow \infty \quad (10c).$$

This function diverges (goes to zero) when $\delta < 0$ ($\delta > 0$).

Since $\chi \rightarrow 0$ (for $\xi^{-1} = 0$) corresponds to $\rho_1 \rightarrow 0$, the perturbation expansion leading to Eq.(9) is valid only for $y = -\ln \chi \ll 1/\epsilon$. Hence, to determine the correct boundary condition as $y \rightarrow \infty$ the solution of Eq.(9) must be matched with the "inner" solution describing the correlator with two points near coincidence. The latter is governed by the equation derived directly from Eq.(3) by expanding in $\rho_1/\rho_2 \ll 1$ instead of ϵ and which is written conveniently in the polar coordinates $|\rho_1/\rho_2|^2 = \xi^{-2} + \chi^2/4$ and $\theta = \arctan(2/\xi\chi)$ (restricting here to $d = 2$ for simplicity). The natural radial variable in this "inner" equation turns out to be $Y = |\rho_1|^\epsilon$. The region of matching with Eq.(10a) corresponds to $1 - Y \ll 1$ and $\theta = 0$. Quite generally the solution near $Y = 1$ behaves as $A + B(1 - Y)^\alpha$ with $\alpha > 0$ required to keep the solution from diverging. In the matching region $Y \approx 1 + \epsilon \ln(\chi/2)$ so that only the constant term, A , must be kept when computing to the leading order in ϵ . Comparing with Eq.(10c) one concludes that matching the "inner" solution is only possible for $\delta = 0$.

For $\delta = 0$ the solution of Eq.(9) is $a(\chi) = \sin(\pi/6 - \chi)/\sin(\pi/6)$ for $0 < \chi < \pi/3$ which is continued over the full range of χ using reflection symmetry and periodicity defined above. One observes that $a(\chi)$ has an apparent $|\chi|$ singularity near $\chi = 0$ (and other points related by symmetry) which is regularized only for $\chi < e^{-1/\epsilon}$ via the crossover to the "inner" solution for nearly coincident points $\rho_1/\rho_2 < e^{-1/\epsilon}$ as discussed above. Note that although to $O(\epsilon)$ there is no correction to the $\lambda = 1$ eigenvalue, the computed eigenfunction is non-trivial: it is a superposition of many $\psi_{1,q}$ modes since $a_q \sim 1/q$ for large q . Also note that the calculations in 2 and 3-dimensions are identical and give the same result: $\delta = 0$.

Thus, the main result of this paper : the scaling exponent of the $n = 3$ structure function behaves as $\lambda = 1$, up to corrections smaller than ϵ^1 . The exponent of the 3-point

correlation function is therefore smaller than the "naive" scaling exponent, equal to $1 + \epsilon$, therefore demonstrating that the behavior of the skewness near the Batchelor limit of the Kraichnan, white velocity, model is anomalous. Dispersion in the presence of a mean gradient has been shown experimentally^{19,20} and numerically^{21,22} to give rise to strong intermittency effects, resulting in a skewness which remains of order 1, independent of the Reynolds number. As it is the case in real flows, it is interesting to notice that even for a white noise velocity field, the anisotropy induced at large scales decays more slowly than predicted by standard phenomenological arguments.^{12,13,23} We conclude by mentioning numerical results demonstrating that a large scale anisotropy, such as a large scale shear, imposed on a turbulent velocity field, may also result in a large anisotropy at small scale²⁴, suggesting also the existence of an anomalous exponent for the $n = 3$ structure function of the velocity field.

Acknowledgements

AP thanks DRET for support under contract # 95-2591A, and EDS was supported by the NSF under contract # DMR9121654

References

- [1] R. H. Kraichnan, Phys. Rev. Lett. **72**, 1016 (1994).
- [2] R. H. Kraichnan, Phys. Fluids **11**, 945 (1968).
- [3] K. Gawędzki and A. Kupiainen, Phys. Rev. Lett. **75**, 3834 (1995).
- [4] M. Chertkov *et al.*, Phys. Rev. **E 52**, 4924 (1995).
- [5] B.I. Shraiman and E.D. Siggia, C. R. Acad. Sci. Paris, **321**, Sér. IIb, 279 (1995).
- [6] B. I. Shraiman and E. D. Siggia, Phys. Rev. **E49**, 2912 (1994).
- [7] R. H. Kraichnan, V. Yakhot and S. Chen, Phys. Rev. Lett. **75**, 240, (1995).
- [8] R. H. Kraichnan, J. Fluid Mech. **64**, 737 (1974).
- [9] B. I. Shraiman and E. D. Siggia, Phys. Rev. Lett, **77**, 2463, (1996).
- [10] A. Pumir, Europh. Lett. **34**, 25 (1996).
- [11] D. Gutman and E. Balkovsky, preprint, chao-dyn/9604003, (1996).
- [12] S. Corrsin, NACA R & M 58B1 (1958).
- [13] J.L. Lumley, Phys. Fluids **10**, 855 (1967).
- [14] R. A. Antonia and C. W. Van Atta, J. Fluid Mech. **84**, 561 (1980).
- [15] G. K. Batchelor, J. Fluid Mech. **5**, 113 (1959).
- [16] See also E.Balkovsky et al., JETP Lett., **61**, 1021 (1995)), who investigated the 4 point correlator.
- [17] L. Landau and E. Lifshitz, *Quantum Mechanics*, Pergamon Press, New-York (1986).
- [18] G. Vilenkin, *Representations of groups and special functions* , Kluwer Acad. Publ., (1991).
- [19] K.R. Sreenivasan, Proc. R. Soc. London **A 434**, 165 (1991).
- [20] C. Tong and Z. Warhaft, Phys. Fluids **6**, 2165 (1994).
- [21] M. Holzer and E.D. Siggia, Phys. Fluids **6**, 1820 (1994).
- [22] A. Pumir, Phys. Fluids **6**, 2118 (1994).
- [23] While this paper was in preparation, we learned of the numerical study of the spectrum of the 3-point Hopf operator for the Kraichnan model by O. Gat, V. S. L’vov,

E. Podivilov and I. Procaccia (submitted to Phys. Rev. Lett.). However their calculation addressed the $l = 0$ angular sector, which by symmetry does not contribute to the 3-d order structure function. It would be interesting to carry out the numerical diagonalization in the $l=1$ sector which could be compared with the anomalous skewness exponents found in Ref. 9 and in the present paper.

[24] A. Pumir and B. I. Shraiman, Phys. Rev. Lett. **75** , 3114 (1995).